

Thalamus as a “hub” to predict outcome after epilepsy surgery

Barbara C. Jobst, MD, Dr
Med
Gregory D. Cascino, MD

Correspondence to
Dr. Jobst:
Barbara.C.Jobst@hitchcock.org

Neurology® 2017;88:1–2

The role of the thalamus in mediating seizure activity has been investigated for decades.¹ In 1952, Penfield² proposed “centrencephalic” interactions as an integral part not only of generalized but also of focal automatisms. The dynamic integration between the centrencephalic system and cerebral cortex was hypothesized as pivotal for normal brain function, as well as a mechanism for seizure propagation. The concept was later hotly debated, and generalized seizures with 3-Hz spike wave discharges were considered mainly centrencephalic.¹

Recently, the thalamus has again gained increased importance in focal epilepsy. Intermittent stimulation of the anterior nucleus of the thalamus decreases the seizure frequency in focal epilepsy.³ Since Penfield’s time, we have made tremendous progress in imaging the brain. However, our concepts of epileptic seizures and epilepsy may not have progressed at the same pace.

In this issue of *Neurology*®, He et al.⁴ present a study using resting-state MRI to predict the outcomes of epilepsy surgery. They apply graph theory to measure the “hubness” of 45 nodes per hemisphere and their integration into the global network. They compare seizure-free with not seizure-free patients after epilepsy surgery and with normal controls. All patients had similar temporal lobectomies. One significant finding is that not seizure-free patients had a greater number of and more important connections in the ipsilateral and contralateral thalamus, while seizure-free patients were equivalent to normal controls. No differences were found in other examined regions, including the contralateral temporal lobe. Nodes in seizure-free and not seizure-free patients differed in the richness of connections (degree centrality) and in the influence of the node on the network (eigenvector centrality) but not in modulating signals between nodes (between centrality). Thalamic nodes played a more important role in network integration in not seizure-free patients compared to seizure-free patients, as measured by global efficiency. This demonstrates that patients are less likely to be seizure-free if the thalamus, ipsilateral and

contralateral, is integrally part of the epileptic network and functions as a major “hub.”

Because graph theory measures are theoretical constructs, the authors confirmed their findings with an additional analysis in the supplementary materials. Graph theory measures do not necessarily reflect anatomic connections. It remains to be discussed which one of the interthalamic structures is most relevant to the maintenance or recurrence of seizures and at least theoretically would be the most advantageous target for stimulation approaches for the treatment of epilepsy.

This work nicely complements animal studies in which thalamocortical connections are necessary to sustain seizure activity in a cortical injury model of epilepsy, a well-defined model of focal epilepsy.⁵ In addition, seizures in the injury model could be controlled by optogenetically modulating thalamocortical pathways in a closed-loop paradigm.⁵ Other rodent studies found that involvement of thalamic structures correlated with seizure severity in a kindling model of seizures.⁶

In the work of He et al., thalamic measures of hubness and network integration predicted outcome after epilepsy with greater accuracy than routine clinical measures (74% vs 56%). Another recent study focused on volumetric and diffusion tensor imaging and correlated it to surgical outcome.⁷ In that study, only pulvinar and dorsomedial thalamic volume compared to volumes of any other structures (including volume of resection) correlated with outcome after surgery.⁷ Thalamic and subcortical structures are involved in secondary generalization of focal seizures.⁸ As a clinical measure, a history of secondary generalization is a negative predictor of outcome of seizures. It is possible that if generalization occurs, larger networks are involved in the maintenance of seizures. Along these lines, resting-state fMRI measures and clinical variables may nicely complement each other.

In addition to clinical measures, the study by He et al. may give us another tool to predict outcome after epilepsy surgery.⁹ It seems that network models of epilepsy increasingly substitute for purely focal models of epilepsy.¹⁰ We could imagine a model of epilepsy in

See page 2285

From the Geisel School of Medicine at Dartmouth (B.C.J.), Dartmouth-Hitchcock Medical Center, Hanover, NH; and Mayo Clinic (G.D.C.), Rochester, MN.

Go to *Neurology*.org for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the editorial.

which seizures would be divided into seizures with larger and smaller networks to predict outcome after epilepsy surgery. In clinical practice, it seems premature not to proceed to surgery if such measures of increased hubness in the thalamus are present. However, additional predictive measures could prove helpful in counseling patients. In the past, MRI was mainly predictive insofar as whether a lesion was present or not; additional resting-state MRI may give us more precise predictive measures.⁹

Furthermore, the He et al. study supports the concept of treating seizures with stimulation of thalamic targets after unsuccessful surgery. In the limited experience with stimulation of the anterior thalamic nucleus after unsuccessful epilepsy surgery, stimulation decreased seizure frequency to an extent similar to that of patients without previous surgery.³

As Penfield² already suggested 65 years ago, the thalamus and thalamocortical connections are of great importance even in focal or temporal lobe epilepsy. Thalamocortical connections play a larger role in the epileptic network than previously assumed, and they are certainly worthy of further investigation.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

Dr. Jobst has contracts as researcher through Dartmouth-Hitchcock with Medtronic, Inc and Neuropace, Inc. She also has research support from the Centers for Disease Control and Prevention, Defense Advanced Research Projects Agency, and National Science Foundation. Dr. Cascino is an associate editor for *Neurology*[®] and has received royalties through Mayo Medical Ventures (Nevro Corp). He has also received honoraria

from the American Academy of Neurology and the American Epilepsy Society. Go to Neurology.org for full disclosures.

REFERENCES

1. Jasper HH. Current evaluation of the concepts of centrencephalic and cortico-reticular seizures. *Electroencephalogr Clin Neurophysiol* 1991;78:2–11.
2. Penfield W. Epileptic automatism and the centrencephalic integrating system. *Res Publ Assoc Res Nerv Ment Dis* 1952;30:513–528.
3. Fisher R, Salanova V, Witt T, et al. Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. *Epilepsia* 2010;51:899–908.
4. He X, Doucet GE, Pustina D, Sperling MR, Sharan AD, Tracy JJ. Presurgical thalamic “hubness” predicts surgical outcome in temporal lobe epilepsy. *Neurology* 2017;88:2285–2293.
5. Paz JT, Davidson TJ, Frechette ES, et al. Closed-loop optogenetic control of thalamus as a tool for interrupting seizures after cortical injury. *Nat Neurosci* 2013;16:64–70.
6. Blumenfeld H, Rivera M, Vasquez JG, et al. Neocortical and thalamic spread of amygdala kindled seizures. *Epilepsia* 2007;48:254–262.
7. Keller SS, Richardson MP, Schoene-Bake JC, et al. Thalamotemporal alteration and postoperative seizures in temporal lobe epilepsy. *Ann Neurol* 2015;77:760–774.
8. Blumenfeld H, Varghese GI, Purcaro MJ, et al. Cortical and subcortical networks in human secondarily generalized tonic-clonic seizures. *Brain* 2009;132:999–1012.
9. Jobst BC, Cascino GD. Resective epilepsy surgery for drug-resistant focal epilepsy: a review. *JAMA* 2015;313:285–293.
10. Khambhati AN, Davis KA, Lucas TH, Litt B, Bassett DS. Virtual cortical resection reveals push-pull network control preceding seizure evolution. *Neuron* 2016;91:1170–1182.

Neurology®

Thalamus as a "hub" to predict outcome after epilepsy surgery

Barbara C. Jobst and Gregory D. Cascino

Neurology published online May 17, 2017

DOI 10.1212/WNL.0000000000004043

This information is current as of May 17, 2017

Updated Information & Services

including high resolution figures, can be found at:

<http://www.neurology.org/content/early/2017/05/17/WNL.0000000000004043.full.html>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://www.neurology.org/misc/about.xhtml#permissions>

Reprints

Information about ordering reprints can be found online:

<http://www.neurology.org/misc/addir.xhtml#reprintsus>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2017 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

